

Left-Right squarks mixings effects in Charged Higgs Bosons decays $H^\pm \rightarrow W^\pm(\gamma, Z)$ in the MSSM

Abdesslam Arhrib^{1,2}, Rachid Benbrik² and Mohamed Chabab²

¹ Département de Mathématiques, Faculté des Sciences et Techniques B.P 416
Tanger, Morocco.

² LPHEA, Département de Physique, Faculté des Sciences-Semlalia, B.P. 2390
Marrakech, Morocco.

Abstract

We study the complete one loop contribution to $H^\pm \rightarrow W^\pm V$, $V = Z, \gamma$, in the Minimal Supersymmetric Standard Model (MSSM). We evaluate the MSSM contributions taking into account $B \rightarrow X_s \gamma$ constraint as well as experimental constraints on the MSSM parameters. In the MSSM, we found that in the intermediate range of $\tan \beta \lesssim 10$ and for large A_t and large μ , where lightest stop becomes very light and hence squarks contribution is not decoupling, the branching ratio of $H^\pm \rightarrow W^\pm Z$ can be of the order 10^{-3} while the branching ratio of $H^\pm \rightarrow W^\pm \gamma$ is of the order 10^{-5} . We also study the effects of the CP violating phases of Soft SUSY parameters and found that they can modify the branching ratio by about one order of magnitude.

1 Introduction

The Standard Model (SM) of electroweak interactions is very successful in explaining all experimental data available till now. The cornerstone of the SM, the electroweak symmetry breaking mechanism, still has to be established and the Higgs boson has to be discovered. The main goals of future colliders such as LHC and ILC is to study the scalar sector of the SM. Moreover, the problematic scalar sector of the SM can be enlarged and some simple extensions such as the Minimal Supersymmetric Standard Model (MSSM) and the Two Higgs Doublet Model (2HDM) [1, 2] are intensively studied. Both in the 2HDM and MSSM the electroweak symmetry breaking is generated by 2 Higgs doublets fields Φ_1 and Φ_2 . After electroweak symmetry breaking we are left with 5 physical Higgs particles (2 charged Higgs $H^\pm$, 2 CP-even H^0, h^0 and one CP-odd A^0). The charged Higgs $H^\pm$, because of its electrical charge, is noticeably different from the other SM or 2HDM/MSSM Higgs particles, its discovery would be a clear evidence of physics beyond the SM. In this

study, our concerns is charged Higgs decays $H^\pm \rightarrow W^\pm V$, $V = Z, \gamma$, we will first review the production mechanisms of charged Higgs.

The charged Higgs can be copiously produced both at hadrons and e^+e^- colliders. In hadronic machines, the charged Higgs bosons can be produced in many channels: *i)* the production of $t\bar{t}$ pairs may offer a source of charged Higgs production. If kinematically allowed $m_{H^\pm} \lesssim m_t$, the top quark can decay to $H^+\bar{b}$, competing with the SM decay $t \rightarrow W^+b$. This mechanism can provide a larger production rate of charged Higgs and offers a much cleaner signature than that of direct production.

ii) single charged Higgs production via $gb \rightarrow tH^-$, $gg \rightarrow t\bar{b}H^-$, $qb \rightarrow q'bH^-$ [3]. *iii)* single charged Higgs production in association with $W^\pm$ gauge boson via $gg \rightarrow W^\pm H^\mp$ or $b\bar{b} \rightarrow W^\pm H^\mp$ [4] and also single charged Higgs production in association with A^0 boson via $qq, gg \rightarrow A^0 H^\mp$ [5]. *iv)* $H^\pm$ pair production through $q\bar{q}$ annihilation [6] or gluon fusion.

At e^+e^- colliders, the simplest way to get a charged Higgs is through $H^\pm$ pair production. Such studies have been already undertaken at tree-level [7] and one-loop orders [8] and shown that e^+e^- machines will offer a clean environment and in that sense a higher mass reach.

Experimentally, the null-searches from L3 collaborations at LEP-II derive the lower limit of about $m_{H^\pm} \gtrsim 80$ GeV [9], a limit which applies to all models (2HDM or MSSM) in which $\text{BR}(H^\pm \rightarrow \tau\nu_\tau) + \text{BR}(H^\pm \rightarrow cs) = 1$. DELPHI has also carried out search for $H^\pm \rightarrow A^0 W^\pm$ ¹ topologies in the context of 2HDM type I and derive the lower limit of about $m_{H^\pm} \gtrsim 76$ GeV [10]. Recently and for relatively small $\tan\beta \lesssim 1$ and for a specific SUSY spectrum, CDF Run II can excluded a charged Higgs mass in the range $80 < m_{H^\pm} < 160$ GeV [14]. While for intermediate range of $\tan\beta$ CDF has no limit. If the charged Higgs decay exclusively to $\bar{\tau}\nu$, the $\text{BR}(t \rightarrow H^+b)$ is constrained to be less than 0.4 at 95%C.L. On the other hand if no assumption is made on charged Higgs decay, the $\text{BR}(t \rightarrow H^+b)$ is constrained to be less than 0.91 at 95%C.L.

At the LHC, the detection of light charged Higgs boson with $m_{H^\pm} \lesssim m_t$ is straightforward from top production followed by the decay $t \rightarrow bH^{+2}$. Such light charged Higgs ($m_{H^\pm} \lesssim m_t$) can be detected also for any $\tan\beta$ in the $\tau\nu$ decay which is indeed the dominant decay mode [15]. However, for heavy charged Higgs masses $m_{H^\pm} \gtrsim m_t$ which decay predominantly to $t\bar{b}$, the search is rather difficult due to large irreducible and reducible backgrounds associated with $H^+ \rightarrow t\bar{b}$ decay. However, it has been demonstrated in [16] that the $H^+ \rightarrow t\bar{b}$ signature can lead to a visible signal at LHC provided that the charged Higgs mass below 600 GeV and $\tan\beta$ is either below $\lesssim 1.5$ or above $\gtrsim 40$. Ref. [17], proposed $H^\pm \rightarrow \tau\nu$ as an alternative decay mode to detect a heavy charged Higgs, even if such decay is suppressed for heavy charged Higgs it has the advantage being more clean than $H^+ \rightarrow t\bar{b}$.

An other alternative discovery channel for heavy charged Higgs is its decay to charged gauge boson and lightest CP-even Higgs: $H^\pm \rightarrow W^\pm h^0$, followed by the dominant decay of h^0 to $b\bar{b}$ [18]. Since the branching ratio of $H^\pm \rightarrow W^\pm h^0$ is suppressed for High $\tan\beta$,

¹Note that in the 2HDM it may be possible that the decay channel $H^\pm \rightarrow W^\pm A^0$ is open and even dominate over $\tau\nu$ mode for $m_{H^\pm} \lesssim m_t$ [11, 12, 13].

²Note that at Tevatron run II, the charged Higgs is also searched in top decay [14].

this channel could lead to charged Higgs discovery only for low $\tan\beta$ where the branching ratio of $H^\pm \rightarrow W^\pm h^0$ is sizeable.

In MSSM, at tree level, the coupling $H^\pm \rightarrow W^\pm \gamma$ is absent because of electromagnetic gauge invariance $U(1)_{\text{em}}$. While the absence of $H^\pm \rightarrow W^\pm Z$ is due to the isospin symmetry of the kinetic Lagrangian of the Higgs fields [19]. Therefore, decays modes like $H^\pm \rightarrow W^\pm \gamma$, $H^\pm \rightarrow W^\pm Z$ are mediated at one loop level and then are expected to be loop suppressed [20, 21, 22, 23, 24]. We emphasize here that it is possible to construct models with an even larger scalar sector than 2 Higgs doublets, one of the most popular being the Higgs Triplet Model (HTM) [25]. A noteworthy difference between 2HDM and HTM is that the HTM contains a tree level $ZW^\pm H^\mp$ coupling.

Motivated by the fact that there is no detailed study about $H^\pm \rightarrow W^\pm V$, $V = Z, \gamma$, in the framework of MSSM in the literature which take into account left-right squarks mixing, $b \rightarrow s\gamma$ and other electroweak and experimental constraints. We would like to reconsider and update the existing works [20, 21, 22, 23, 24] on the charged Higgs boson decays into a pair of gauge boson: $H^\pm \rightarrow W^\pm \gamma, W^\pm Z$ both in 2HDM and MSSM with and without CP violating phases. Although these decays are rare processes, loop or/and threshold effects can give a substantial effect. Moreover, once worked out, any experimental deviation from the results within such a model should bring some fruitful information on the new physics and allow to distinguish between models. We would like to mention also that, those channels have a very clear signature and might emerge easily at future colliders. For instance, if $H^\pm \rightarrow W^\pm Z$ is enhanced enough, this decay may lead to three leptons final state if both W and Z decay leptonically and that would be the corresponding golden mode for charged Higgs boson.

Charged Higgs decays: $H^\pm \rightarrow W^\pm \gamma, W^\pm Z$, have received much more attention in the literature. $H^\pm \rightarrow W^\pm Z$ has been studied first in the MSSM in [20]. Ref. [21] has considered both $H^\pm \rightarrow W^\pm \gamma$ and $H^\pm \rightarrow W^\pm Z$ in the MSSM and show that the rate of $H^\pm \rightarrow W^\pm \gamma$ is very small while the rate of $H^\pm \rightarrow W^\pm Z$ can be enhanced by heavy fermions particles in the loops. The fourth generation contribution was given as an example. Although the squarks contribution has been considered in Ref. [21], Left-Right squarks mixing which could give substantial enhancement has been neglected. In contrast to Ref. [21] which argue that the squarks contributions decouple, we will show that there is non-decoupling effects originating from squarks contributions at large A_t and large μ limit. $H^\pm \rightarrow W^\pm \gamma$ was also studied in [22] within the MSSM, but the pure SUSY contribution from charginos, neutralinos and squarks has been neglected. Later on, Ref. [23] studied the possibility of enhancing $H^\pm \rightarrow W^\pm Z$ by the non-decoupling effect of the heavy Higgs bosons in the context of 2HDM, substantial enhancement was found [23]. Recently, $H^\pm \rightarrow W^\pm \gamma$ was also studied in 2HDM type II [24]. All the above studies has been carried out either in unitary gauge [20, 21] or in the nonlinear R_ξ -gauge [24]. The analysis of [22] and [23] have been performed in 'tHooft-Feynman gauge without any renormalization scheme. It has been checked in [22, 23] that the sum of all Feynman diagrams: vertex, tadpoles and vector boson-scalar mixing turns out to be Ultra-Violet finite.

In the present study, we will still use 'tHooft-Feynman gauge to do the computation. However, the amplitudes of $H^\pm \rightarrow W^\pm \gamma$ and $H^\pm \rightarrow W^\pm Z$ are absent at the tree level,

complications like tadpoles contributions and vector boson–scalar mixing require a careful treatment of renormalization. We adopt hereafter the on-shell renormalization scheme developed in [26].

The paper is organized as follows. In section II, we describe our calculations and the one-loop renormalization scheme we will use for $H^\pm \rightarrow W^\pm Z$ and $H^\pm \rightarrow W^\pm \gamma$. In Section III, we present our numerical results and discussions, and section VI contains our conclusions.

2 Charged Higgs decay: $H^\pm \rightarrow W^\pm V$

As we have seen in the previous section, in MSSM, at tree level, the coupling $H^\pm \rightarrow W^\pm \gamma$ and $H^\pm \rightarrow W^\pm Z$ do not exist. They are generated at one loop level and then are expected to be loop suppressed [20, 21, 22, 23, 24]. Hereafter, we will give the general structure of such one loop couplings and discuss the renormalization scheme introduced to deal with tadpoles and vector boson scalar boson mixing.

2.1 One loop amplitude $H^\pm \rightarrow W^\pm V$

The amplitude $\mathcal{M}$ for a scalar decaying to two gauges bosons V_1 and V_2 can be written as

$$\mathcal{M} = \frac{g^3 \epsilon_{V_1}^{\mu*} \epsilon_{V_2}^{\nu*}}{16\pi^2 m_W} \mathcal{M}_{\mu\nu} \quad (1)$$

where ϵ_{V_i} are the polarization vectors of the V_i .

According to Lorenz invariance, the general structure of the one loop amplitude $\mathcal{M}_{\mu\nu}$ of $S \rightarrow V_1^\mu V_2^\nu$ decay, if CP is conserved, is

$$\mathcal{M}_{\mu\nu}(S \rightarrow W^\mu V^\nu) = \mathcal{F}_1 g_{\mu\nu} + \mathcal{F}_2 p_{1\mu} p_{2\nu} + \mathcal{F}_3 i \epsilon_{\mu\nu\rho\sigma} p_1^\rho p_2^\sigma \quad (2)$$

where $p_{1,2}$ are the momentum of V_1, V_2 vector bosons, $\mathcal{F}_{1,2,3}$ are form factors, and $\epsilon_{\mu\nu\rho\sigma}$ is the totally antisymmetric tensor. The form factor $\mathcal{F}_1$ has dimension 2 while the other are dimensionless.

For $H^\pm \rightarrow W^\pm \gamma$, electromagnetic gauge invariance implies that $\mathcal{F}_1 = 1/2(m_W^2 - m_{H^\pm}^2) \mathcal{F}_2$ [21]. This means that only $\mathcal{F}_2$ and $\mathcal{F}_3$ will contribute to the decay $H^\pm \rightarrow W^\pm \gamma$. In case of $H^\pm \rightarrow W^\pm Z$, there is no such constraint on form factors.

In terms of an effective Lagrangian analysis, from gauge invariance requirement we can write:

$$\mathcal{L}_{eff} = g_1 H^\pm W_\mu^\mp V^\mu + g_2 H^\pm F_V^{\mu\nu} F_{W\mu\nu} + i g_3 \epsilon_{\mu\nu\rho\sigma} H^\pm F_V^{\mu\nu} F^{W\rho\sigma} + h.c \quad (3)$$

the first operator $H^\pm W_\mu^\mp V^\mu$ is dimension three and the last two operators $H^\pm F_V^{\mu\nu} F_{W\mu\nu}$ and $\epsilon_{\mu\nu\rho\sigma} H^\pm F_V^{\mu\nu} F^{W\rho\sigma}$ are dimension five. One conclude that $g_{2,3}$ (resp. g_1) must be of the form $g(R)/M$ (resp $Mg(R)$) with M a heavy scale in MSSM, $g(R)$ a dimensionless function and R is a ratio of some internal masses of the model under studies. Therefore, it is expected that in case of $H^\pm \rightarrow W^\pm Z$ decay, $\mathcal{F}_1$ will grow quadratically with internal

top quark mass while $\mathcal{F}_{2,3}$ will have only logarithmic dependence [21]³. A contrario, for $H^\pm \rightarrow W^\pm \gamma$ decay, the electromagnetic gauge invariance relates $\mathcal{F}_1$ and $\mathcal{F}_2$ and then the amplitude of $H^\pm \rightarrow W^\pm \gamma$ will not grows quadratically with internal masses. One expect that the decay $H^\pm \rightarrow W^\pm \gamma$ is less enhanced compared to $H^\pm \rightarrow W^\pm Z$.

2.2 On-shell renormalization

We have evaluated the one-loop induced process $H^\pm \rightarrow W^\pm V$ in the 'tHooft-Feynman gauge using dimensional regularization. Since we are interested in the charged Higgs decay to SM particles like $W^\pm Z$ and $W^\pm \gamma$, at one-loop level, dimensional regularization will give the same result as dimensional reduction. In fact, we have checked numerically that the dimensional regularization and dimensional reduction gives the same result.

The typical Feynman diagrams that contribute to $H^\pm \rightarrow W^\pm V$ are depicted in Fig. 1. Those diagrams contains vertex diagrams (Fig. 1.1 $\rightarrow$ 1.11), $W^\pm$ - $H^\pm$ mixing (Fig. 1.12 $\rightarrow$ 1.14), $H^\pm$ - $G^\pm$ mixing (Fig. 1.15 $\rightarrow$ 1.17) and $H^\pm$ - $W^\pm$ mixing (Fig. 1.18 $\rightarrow$ 1.20).

Note that the mixing $H^\pm$ - $W^\pm$ (Fig. 1.12, 1.13, 1.14) vanishes for an on-shell transverse W gauge boson. There is no contribution from the $W^\pm$ - $G^\mp$ mixing because $\gamma G^\pm H^\mp$ and $Z G^\pm H^\mp$ vertices are absent at the tree level. All the Feynman diagrams have been generated and computed using FeynArts and FormCalc [27] packages. We also used the fortran FF-package [28] in the numerical analysis.

Although the amplitude for our process is absent at the tree level, complications like tadpole contributions and vector boson-scalar mixing require a careful treatment of renormalization. We adopt, hereafter, the on-shell renormalization scheme of [29], for the Higgs sector, which is an extension of the on-shell scheme in [30]. In this scheme, field renormalization is performed in the manifest-symmetric version of the Lagrangian. A field renormalization constant $Z_{\Phi_{1,2}}$ is assigned to each Higgs doublet $\Phi_{1,2}$. Following the same approach adopted in [26], the Higgs fields and vacuum expectation values v_i are renormalized as follows:

$$\begin{aligned}\Phi_i &\rightarrow (Z_{\Phi_i})^{1/2} \Phi_i \\ v_i &\rightarrow (Z_{\Phi_i})^{1/2} (v_i - \delta v_i).\end{aligned}\tag{4}$$

With these substitutions in the scalar covariant derivative Lagrangian of the Higgs fields (in the convention of [1]), followed by expanding the renormalization constants $Z_i = 1 + \delta Z_i$ to the one-loop order, we obtain all the counter-terms relevant for our process:

$$\delta[W_\nu^\pm H^\mp] = k^\mu \Delta \tag{5}$$

$$\delta[A_\nu W_\mu^\pm H^\mp] = -ie g_{\mu\nu} \Delta \tag{6}$$

$$\delta[Z_\nu W_\mu^\pm H^\mp] = -ie g_{\mu\nu} \frac{s_W}{c_W} \Delta \tag{7}$$

³As it has been shown in Ref. [21], the top quark contribution does not decouple while squarks contributions does




Figure 1: Generic contributions to $H^\pm \rightarrow W^\pm V$

where k denotes the momentum of the incoming $W^\pm$ and

$$\Delta = \frac{\sin 2\beta}{2} m_W \left[\frac{\delta v_2}{v_2} - \frac{\delta v_1}{v_1} + \delta Z_{\Phi_1} - \delta Z_{\Phi_2} \right]. \quad (8)$$




Figure 2: Generic contributions to $H^\pm \rightarrow W^\pm$ and $H^\pm \rightarrow G^\pm$ mixing as well as counter-terms needed.

Denoting the one particle irreducible (1PI) two point function for $W^\pm H^\pm$ (resp $G^\pm H^\pm$) mixing by $\pm i k_\mu \Sigma_{W^\pm H^\pm}(k^2)$ (resp $i \Sigma_{G^\pm H^\pm}(k^2)$) where k is the momentum of the incoming $W^\pm$ (resp $G^\pm$), and $H^\pm$ is outgoing. The renormalized mixing will be denoted by $\hat{\Sigma}$.

In the on-shell scheme, we will use the following renormalization conditions:

- The renormalized tadpoles, i.e. the sum of tadpole diagrams $T_{h,H}$ and tadpole counter-terms $\delta_{h,H}$ vanish:

$$T_h + \delta t_h = 0, \quad T_H + \delta t_H = 0.$$

These conditions guarantee that $v_{1,2}$ appearing in the renormalized Lagrangian $\mathcal{L}_{\mathcal{R}}$ are located at the minimum of the one-loop potential.

- The real part of the renormalized non-diagonal self-energy $\hat{\Sigma}_{H^\pm W^\pm}(k^2)$ vanishes for an on-shell charged Higgs boson:

$$\Re \hat{\Sigma}_{H^\pm W^\pm}(m_{H^\pm}^2) = 0 \quad (9)$$

This renormalization condition determines the term Δ to be

$$\Delta = \Re \Sigma_{H^\pm W^\pm}(m_{H^\pm}^2) \quad (10)$$

and consequently $\delta[A_\nu W_\mu^\pm H^\mp]$ and $\delta[Z_\nu W_\mu^\pm H^\mp]$ are also fixed.

The last renormalization condition is sufficient to discard the real part of the $H^\pm$ - $G^\pm$ mixing contribution as well. Indeed, using the Slavnov–Taylor identity [31]

$$k^2 \Sigma_{H^\pm W^\pm}(k^2) - m_W \Sigma_{H^\pm G^\pm}(k^2) = 0 \quad \text{at} \quad k^2 = m_{H^\pm}^2 \quad (11)$$

which is valid also for the renormalized quantities together with eq. (9), it follows that

$$\Re \hat{\Sigma}_{H^\pm G^\pm}(m_{H^\pm}^2) = 0. \quad (12)$$

In particular, the Feynman diagrams depicted in Fig. 1.9 will not contribute with the above renormalization conditions, being purely real valued.

To make the amplitude of Fig.1 Ultra-Violet finite we need to add the following counter-terms: counter-terms for $\gamma W^\pm H^\mp$ and $Z W^\pm H^\mp$ vertices Fig.2.2a, a counter-term for the $W^\pm$ - $H^\mp$ mixing Fig.2.2b, 2.2d, and a counter-term for the $G^\pm$ - $H^\mp$ mixing Fig. 2.2c.

3 Numerics and discussions

In our numerical evaluations, we use the following experimental input quantities [32]: $\alpha^{-1} = 129$, m_Z , m_W , m_t , $m_b = 91.1875, 80.45, 174.3, 4.7$ GeV. In the MSSM, we specify the free parameters that will be used as follow: i) The MSSM Higgs sector is parameterized by the CP-odd mass m_{A^0} and $\tan \beta$, taking into account one-loop radiative corrections from [33], and we assume $\tan \beta \gtrsim 3$. ii) The chargino–neutralino sector can be parameterized by the gaugino-mass terms M_1 , M_2 , and the Higgsino-mass term μ . For simplification GUT relation $M_1 \approx M_2/2$ is assumed. iii) Sfermions are characterized by a common soft-breaking sfermion mass $M_{SUSY} \equiv \widetilde{M}_L = \widetilde{M}_R$, μ the parameter and the soft trilinear couplings for third generation scalar fermions $A_{t,b,\tau}$. For simplicity, we will take $A_t = A_b = A_\tau$.

When varying the MSSM parameters, we take into account also the following constraints: i) The extra contributions to the $\delta\rho$ parameter from the Higgs scalars should not exceed the current limits from precision measurements [32]: $|\delta\rho| \lesssim 0.003$. ii) $b \rightarrow s\gamma$ constraint. The present world average for inclusive $b \rightarrow s\gamma$ rate is [32] $\mathcal{B}(B \rightarrow X_s \gamma) = (3.3 \pm 0.4) \times 10^{-4}$. We keep the $B \rightarrow X_s \gamma$ branching ratio in the 3σ range of $(2.1\text{--}4.5) \times 10^{-4}$. The SM part of $B \rightarrow X_s \gamma$ is calculated up to NLO using the expression given in [34]. While for the MSSM part, the Wilson coefficient C_7 and C_8 are included at LO in the framework of MSSM with CKM as the only source of flavor violation and are taken from [35]. iii) We will assume that all SUSY particles Sfermions and charginos are heavier than about 100 GeV; for the light CP even Higgs we assume $m_{h^0} \gtrsim 98$ GeV and $\tan \beta \gtrsim 3$ [36].

As the experimental bound on m_{h^0} is concerned, care has to be taken. Since we are using only one-loop approximation for the Higgs spectrum, and as it is known, higher order corrections [37] may reduce the light CP-even Higgs mass in some cases. It may be possible




Figure 3: Branching ratios of $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) as a function of $m_{H^\pm}$ in the MSSM and 2HDM for $M_{SUSY} = 500$ GeV, $M_2 = 175$ GeV, $\mu = -1.4$ TeV and $A_t = A_b = A_\tau = -\mu$ for various values of $\tan \beta$.

that some parameter space points, shown in this analysis, which survive to the experimental limit $m_{h^0} \gtrsim 98$ GeV with one loop calculation may disappear once the higher order correction to the Higgs spectrum are included.

The total width of the charged Higgs is computed at tree level from [2] without any QCD improvement for its fermionic decays $H^\pm \rightarrow \bar{f}f'$. The SUSY channels like $H^+ \rightarrow \tilde{f}_i \tilde{f}'_j$ and $H^+ \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^+$ are included when kinematically allowed. In Fig. 3, we show branching ratio of $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) as a function of charged Higgs mass for $\tan \beta = 16$ and 25. In those plots, we have shown both the pure 2HDM⁴ and the full MSSM contribution. As it can be seen from those plots, both for $H^\pm \rightarrow W^\pm Z$ and $H^\pm \rightarrow W^\pm \gamma$ the 2HDM contribution is rather small. Once we include the SUSY particles, we can see that the Branching fraction get enhanced and can reach 10^{-3} in case of $H^\pm \rightarrow W^\pm Z$ and 10^{-5} in case of $H^\pm \rightarrow W^\pm \gamma$. The source of this enhancement is mainly due to the presence of scalar fermion contribution in the loop which are amplified by threshold effects from the opening of the decay $H^\pm \rightarrow \tilde{t}_i \tilde{b}_j^*$. It turns out that the contribution of charginos neutralinos loops does not enhance the Branching fraction significantly as compared to scalar fermions loops. The plots also show that, the branching fraction is more important for intermediate $\tan \beta = 16$ and is slightly reduced for larger $\tan \beta = 25$.

This $\tan \beta$ dependence is shown in Fig. 4 both for $H^\pm \rightarrow W^\pm Z$ and $H^\pm \rightarrow W^\pm \gamma$ for three representative values of A_t . It is obvious that the smallest is $\tan \beta$ the largest is the branching fraction. Increasing $\tan \beta$ from 5 to about 40 can reduce the branching fraction by about one or two order of magnitude. As one can see from those figures, the plots stops for $\tan \beta \approx 24$ for $A_t = 500$ GeV, this is due to $b \rightarrow s \gamma$ constraint. For $A_t = 1400$ GeV, only $\tan \beta \in [16, 26]$ is allowed, the reason is that for $\tan \beta \lesssim 16$ the light stop $\tilde{t}_1$ becomes

⁴Pure 2HDM means that we include just the 2HDM part of the MSSM that contributes here in the loop, i.e only SM fermion, gauge bosons and Higgs bosons with MSSM sum rules for the Higgs sector.




Figure 4: Branching ratios for $H^\pm \rightarrow W^\pm Z$ (left), $H^\pm \rightarrow W^\pm \gamma$ (right) as a function of $\tan\beta$ in the MSSM with $M_{SUSY} = 500$ GeV, $M_2 = 200$ GeV, $m_{H^\pm} = 500$ GeV, $A_t = A_b = A_\tau = -\mu$ for various values of A_t .




Figure 5: Scatter plot for branching ratios of $H^\pm \rightarrow W^\pm Z$ (left), $H^\pm \rightarrow W^\pm \gamma$ (right) in the $(m_{H^\pm}, \tan\beta)$ plane in the MSSM for $M_{SUSY} = 1$ TeV, $M_2 = 175$ GeV, $A_{t,b,\tau} = M_{SUSY}$ and $\mu = -1$ TeV.




Figure 6: Branching ratios for $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) as a function of A_t in the MSSM with $M_{SUSY} = 500$ GeV, $M_2 = 200$ GeV, $m_{H^\pm} = 500$ GeV, $A_t = A_b = A_\tau = -\mu$ and -2 TeV $< \mu < -0.1$ TeV for various values of $\tan \beta$.

lighter than the experimental limit of 98 GeV and for $\tan \beta \gtrsim 26$, the experimental limit on the light CP even Higgs h^0 is violated. As indicated above, we assume a conservative limit of $m_{h^0} \gtrsim 98$ GeV rather than $m_{h^0} \gtrsim 114$ GeV which should be used in the case of decoupling limit where ZZh^0 coupling mimic the SM one.

We also show a scatter plot Fig. 5 for $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) in $(m_{H^\pm}, \tan \beta)$ plane for $A_t = -\mu = 1$ TeV, $M_{SUSY} = A_t$ and $M_2 = 175$ GeV. As it can be seen from Fig 5 there is only a small area for $\tan \beta \lesssim 10$ where the branching ratio of $H^\pm \rightarrow W^\pm Z$ can be in the range 10^{-5} – 10^{-3} .

We now illustrate in Fig. 6 the branching fraction of $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) as a function of $A_t = A_b = A_\tau = -\mu$ for $M_{SUSY} = 500$ GeV and $M_2 = 200$ GeV. Since $b \rightarrow s\gamma$ favor A_t and μ to have opposite sign, we fix $\mu = -A_t$ and in this sense also μ is varied when A_t is varied. Both for $H^\pm \rightarrow W^\pm Z$ and $H^\pm \rightarrow W^\pm \gamma$, the chargino-neutralino contribution which is rather small decrease with $\mu = -A_t$, the largest is A_t the smallest is chargino-neutralino contribution. As one can see from those figures, the plots stops for $A_t = 1.1$ TeV and $\tan \beta = 3$ because for larger A_t $\delta\rho$ constraint is violated. For $\tan \beta = 1$ and 20, the plots stops for the same reason.

In case of $H^\pm \rightarrow W^\pm Z$, for $A_t \lesssim 1$ TeV it is the pure 2HDM contribution which dominate and that is why it is almost independent of A_t while for large A_t the branching ratio increase with A_t . It is clear that the largest is A_t the largest is the branching ratio which can be of the order of 10^{-3} for $H^\pm \rightarrow W^\pm Z$ with $\tan \beta = 10$. As we know from $h^0 \rightarrow \gamma\gamma$ and $h^0 \rightarrow \gamma Z$ in MSSM [38], the squarks contributions decouple except in the light stop mass and large A_t limit [38]. In $H^\pm \rightarrow W^\pm V$ case, the same situation happen. As we can see from Fig. 6 (left), for intermediate A_t , $300 < A_t < 1000$ GeV, the squarks are rather heavy and hence their contributions is small compared to 2HDM one. While for large A_t the stop becomes very light $\lesssim 200$ GeV and hence enhance $H^\pm \rightarrow W^\pm V$ width. Of




Figure 7: Branching ratios for $H^\pm \rightarrow W^\pm Z$ (left) and $H^\pm \rightarrow W^\pm \gamma$ (right) in the MSSM as a function of $Arg(A_t)$: $M_{SUSY} = 500$ GeV, $M_2 = 150$ GeV, $m_{H^\pm} = 500$ GeV, $A_t = A_b = A_\tau = -\mu = 1$ TeV and for various values of $\tan \beta$.

course this enhancement is also amplified by $H^\pm \tilde{t}_{L,R} \tilde{b}_{R,L}^*$ and $H^\pm \tilde{\tau}_{L,R} \tilde{\nu}_{\tau L}^*$ couplings which are directly proportional to $A_{t,b,\tau}$. In case of $H^\pm \rightarrow W^\pm \gamma$ decay, the pure 2HDM and sfermions contribution are of comparable size, the branching ratio increases with A_t .

We have also studied the effect of the MSSM CP violating phases on charged Higgs decays. Similar study has been done for the single charged Higgs production at hadron collider [39]. It is well known that the presence of large SUSY CP violating phases can give contributions to electric dipole moments of the electron and neutron (EDM) which exceed the experimental upper bounds. In a variety of SUSY models such phases turn out to be severely constrained by such constraints i.e. $Arg(\mu) < (10^{-2})$ for a SUSY mass scale of the order of few hundred GeV [40]. For $H^\pm \rightarrow W^\pm Z$ and $H^\pm \rightarrow W^\pm \gamma$ decays which are sensitive to MSSM CP violating phases through squarks and charginos-neutralinos contributions, it turns out that the effect of MSSM CP violating phases is important and can enhance the rate by about one order of magnitude. For illustration we show in Fig. 7 the effect of $A_{t,b,\tau}$ CP violating phases for $M_{SUSY} = 500$ GeV, $A_{t,b,\tau} = -\mu = 1$ TeV and $M_2 = 150$ GeV. For simplicity, we assume that μ is real. As it is clear, the CP phase of $A_{t,b,\tau}$ can enhance the rates of both $H^\pm \rightarrow W^\pm Z, \gamma$ by more than an order of magnitude. The observed cuts in the plot are due to $b \rightarrow s \gamma$ constraint. The CP violating phases can lead to CP-violating rate asymmetry of $H^\pm$ decays, those issues are going to be addressed in an incoming paper [41].

4 Conclusion

In the framework of MSSM we have studied charged Higgs decays into a pair of gauge bosons namely: $H^\pm \rightarrow W^\pm Z$ and $H^\pm \rightarrow W^\pm \gamma$. In the MSSM we have also studied the effects of MSSM CP violating phases. In contrast to previous studies, we have performed

the calculation in the 'tHooft-Feynman gauge and used a renormalization prescription to deal with tadpoles, $W^\pm-H^\pm$ and $G^\pm-H^\pm$ mixing. The study has been carried out taking into account the experimental constraint on the ρ parameter, $b \rightarrow s\gamma$ constraint. Numerical results for the branching ratios have been presented. In the MSSM, we have shown that the branching ratio of $H^\pm \rightarrow W^\pm Z$ can reach 10^{-3} in some cases while $H^\pm \rightarrow W^\pm \gamma$ never exceed 10^{-5} . The effect of MSSM CP violating phases is also found to be important.

Those Branching ratio of the order 10^{-3} might provide an opportunity to search for a charged Higgs boson at the LHC through $H^\pm \rightarrow W^\pm Z$.

At the end, we would like to mention that some effects shown in this study may be ruled out by the experimental bound on the light CP-even mass if we take into account 2-loop radiative corrections on the Higgs spectrum which have the tendency to reduce the light CP-even mass in some cases. On the other hand, the inclusion of high effects for $H^\pm \rightarrow W^\pm V$ like bottom and top quarks mass running as well as $\tan\beta$ resummation could affect the rate of $H^\pm \rightarrow W^\pm V$.

Acknowledgements This work is supported by PROTARS-III D16/04.

References

- [1] J.F. Gunion, H.E. Haber, G.L. Kane and S. Dawson, *The Higgs Hunter's Guide* (Addison-Wesley, Reading, 1990).
- [2] A. Djouadi, arXiv:hep-ph/0503173.
- [3] V. D. Barger, R. J. N. Phillips and D. P. Roy, Phys. Lett. B **324**, 236 (1994); J. F. Gunion, H. E. Haber, F. E. Paige, W. K. Tung and S. S. D. Willenbrock, Nucl. Phys. B **294**, 621 (1987). R. M. Barnett, H. E. Haber and D. E. Soper, Nucl. Phys. B **306**, 697 (1988). J. L. Diaz-Cruz and O. A. Sampayo, Phys. Rev. D **50**, 6820 (1994). S. Moretti and K. Odagiri, Phys. Rev. D **55**, 5627 (1997).
- [4] D. A. Dicus, J. L. Hewett, C. Kao and T. G. Rizzo, Phys. Rev. D **40**, 787 (1989). A. A. Barrientos Bendezu and B. A. Kniehl, Phys. Rev. D **59**, 015009 (1999); A. A. Barrientos Bendezu and B. A. Kniehl, Phys. Rev. D **63**, 015009 (2001); W. Hollik and S. h. Zhu, Phys. Rev. D **65**, 075015 (2002); O. Brein, W. Hollik and S. Kanemura, Phys. Rev. D **63**, 095001 (2001); E. Asakawa, O. Brein and S. Kanemura, Phys. Rev. D **72** (2005) 055017.
- [5] Q. H. Cao, S. Kanemura and C. P. Yuan, Phys. Rev. D **69**, 075008 (2004).
- [6] A. C. Bawa, C. S. Kim and A. D. Martin, Z. Phys. C **47**, 75 (1990). A. A. Barrientos Bendezu and B. A. Kniehl, Nucl. Phys. B **568**, 305 (2000); A. Krause, T. Plehn, M. Spira and P. M. Zerwas, Nucl. Phys. B **519**, 85 (1998); O. Brein and W. Hollik, Eur. Phys. J. C **13**, 175 (2000).
- [7] S. Komamiya, Phys. Rev. D **38**, 2158 (1988). A. Djouadi, J. Kalinowski and P. M. Zerwas, Z. Phys. C **57**, 569 (1993).

- [8] A. Arhrib and G. Moultaka, Nucl. Phys. B **558**, 3 (1999), ibid A. Arhrib, M. Capdequi Peyranere and G. Moultaka, Phys. Lett. B **341**, 313 (1995); J. Guasch, W. Hollik and A. Kraft, Nucl. Phys. B **596** (2001) 66. E. Asakawa, O. Brein and S. Kanemura, arXiv:hep-ph/0508226. O. Brein, T. Hahn, S. Heinemeyer and G. Weiglein, arXiv:hep-ph/0402053.
- [9] P. Achard *et al.* [L3 Collaboration], Phys. Lett. B **575**, 208 (2003).
- [10] J. Abdallah *et al.* [DELPHI Collaboration], Eur. Phys. J. C **34**, 399 (2004).
- [11] F. M. Borzumati and A. Djouadi, Phys. Lett. B **549**, 170 (2002).
- [12] A. G. Akeroyd, Nucl. Phys. B **544**, 557 (1999).
- [13] A. G. Akeroyd, A. Arhrib and E. Naimi, Eur. Phys. J. C **20**, 51 (2001); A. G. Akeroyd, A. Arhrib and E. M. Naimi, Eur. Phys. J. C **12**, 451 (2000).
- [14] A. Abulencia *et al.* [CDF Collaboration], arXiv:hep-ex/0510065; F. Abe *et al.* [CDF Collaboration], Phys. Rev. Lett. **79**, 357 (1997).
- [15] S. Raychaudhuri and D. P. Roy, Phys. Rev. D **53**, 4902 (1996);
- [16] V. D. Barger, R. J. N. Phillips and D. P. Roy, Phys. Lett. B **324**, 236 (1994); J. F. Gunion, Phys. Lett. B **322**, 125 (1994); D. J. Miller, S. Moretti, D. P. Roy and W. J. Stirling, Phys. Rev. D **61**, 055011 (2000); S. Moretti and D. P. Roy, Phys. Lett. B **470**, 209 (1999);
- [17] K. Odagiri, arXiv:hep-ph/9901432; S. Raychaudhuri and D. P. Roy, Phys. Rev. D **53**, 4902 (1996).
- [18] M. Drees, M. Guchait and D. P. Roy, Phys. Lett. B **471**, 39 (1999); S. Moretti, Phys. Lett. B **481**, 49 (2000).
- [19] J. A. Grifols and A. Mendez, Phys. Rev. D **22**, 1725 (1980).
- [20] A. Mendez and A. Pomarol, Nucl. Phys. B **349**, 369 (1991).
- [21] M. Capdequi Peyranere, H. E. Haber and P. Irulegui, Phys. Rev. D **44**, 191 (1991).
- [22] S. Raychaudhuri and A. Raychaudhuri, Phys. Rev. D **50** (1994) 412, ibid Phys. Lett. B **297**, 159 (1992).
- [23] S. Kanemura, Phys. Rev. D **61**, 095001 (2000).
- [24] J. Hernandez-Sanchez, M. A. Perez, G. Tavares-Velasco and J. J. Toscano, Phys. Rev. D **69**, 095008 (2004).
- [25] J. F. Gunion, R. Vega and J. Wudka, Phys. Rev. D **42**, 1673 (1990).

- [26] A. Arhrib, M. Capdequi Peyranere, W. Hollik and G. Moultaka, Nucl. Phys. B **581**, 34 (2000); H. E. Logan and S. f. Su, Phys. Rev. D **66**, 035001 (2002).
- [27] T. Hahn, Comput. Phys. Commun. **140**, 418 (2001); T. Hahn, C. Schappacher, Comput. Phys. Commun. **143**, 54 (2002); T. Hahn, M. Perez-Victoria, Comput. Phys. Commun. **118**, 153 (1999); J. Küblbeck, M. Böhm, A. Denner, Comput. Phys. Commun. **60**, 165 (1990);
- [28] G. J. van Oldenborgh, Comput. Phys. Commun. **66**, 1 (1991); T. Hahn, Acta Phys. Polon. B **30**, 3469 (1999)
- [29] A. Dabelstein, Z. Phys. C **67**, 495 (1995)
- [30] W. F. Hollik, Fortsch. Phys. **38**, 165 (1990).
- [31] J. A. Coarasa, D. Garcia, J. Guasch, R. A. Jimenez and J. Sola, Eur. Phys. J. C **2**, 373 (1998); M. Capdequi Peyranère, Int. J. Mod. Phys. A **14**, 429 (1999).
- [32] S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592** (2004) 1;
- [33] J. F. Gunion and A. Turski, Phys. Rev. D **39**, 2701 (1989). M. S. Berger, Phys. Rev. D **41**, 225 (1990). Y. Okada, M. Yamaguchi and T. Yanagida, Prog. Theor. Phys. **85**, 1 (1991).
- [34] A. L. Kagan and M. Neubert, Phys. Rev. D **58**, 094012 (1998).
- [35] M. Aoki, G. C. Cho and N. Oshimo, Nucl. Phys. B **554**, 50 (1999)
- [36] ALEPH Collaboration, arXiv:hep-ex/0602042.
- [37] S. Heinemeyer, W. Hollik and G. Weiglein, Phys. Lett. B **455**, 179 (1999), S. Heinemeyer, W. Hollik and G. Weiglein, Eur. Phys. J. C **9**, 343 (1999)
- [38] A. Djouadi, V. Driesen, W. Hollik and J. I. Illana, Eur. Phys. J. C **1**, 149 (1998); A. Djouadi, V. Driesen, W. Hollik and A. Kraft, Eur. Phys. J. C **1**, 163 (1998)
- [39] See J. Williams in S. Kraml *et al.*, “Workshop on CP studies and non-standard Higgs physics,” arXiv:hep-ph/0608079.
- [40] P. Nath, Phys. Rev. Lett. **66** (1991) 2565; Y. Kizukuri and N. Oshimo, Phys. Rev. D **46**, 3025 (1992). S. Pokorski, J. Rosiek and C. A. Savoy, Nucl. Phys. B **570**, 81 (2000).
- [41] A. Arhrib, R. Benbrik, W. T. Chang, W.-Y. Keung and T. C. Yuan, work in progress.